

BEST AVAILABLE COPY

OXFORD DICTIONARY OF
BIOCHEMISTRY AND
MOLECULAR BIOLOGY

General Editors **A. D. Smith** (Managing Editor) University College London
S. P. Datta University College London
G. Howard Smith University College London
P. N. Campbell University College London
R. Bentley University of Pittsburgh
H. A. McKenzie University of New South Wales, Australian
Defence Force Academy, and the Australian
National University, Canberra

Subject Editors **D. A. Bender** University College London
A. J. Carozzi University of Queensland
T. W. Goodwin FRS University of Liverpool
J. H. Parish University of Leeds
S. C. Stanford University College London

Oxford University Press
Oxford New York Tokyo
1997

Oxford University Press, Great Clarendon Street, Oxford OX2 6DP

Oxford New York

Athens Auckland Bangkok Bogota Bombay Buenos Aires
Calcutta Cape Town Dar es Salaam Delhi Florence Hong Kong
Istanbul Karachi Kuala Lumpur Madras Madrid Melbourne
Mexico City Nairobi Paris Singapore Taipei Tokyo Toronto

and associated companies in
Berlin Ibadan

Oxford is a trade mark of Oxford University Press

Published in the United States
by Oxford University Press, Inc., New York

© The General Editors, 1997

The authors have asserted their right under the Copyright, Designs and Patents Act, 1988,
to be identified as authors of this work.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without the prior permission in writing of Oxford University Press. Within the UK, exceptions are allowed in respect of any fair dealing for the purpose of research or private study, or criticism or review, as permitted under the Copyright, Designs and Patents Act, 1988, or in the case of reprographic reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency. Enquiries concerning reproduction outside those terms and in other countries should be sent to the Rights Department, Oxford University Press, at the address above.

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form of binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data
(Data applied for)

ISBN 0 19 854768 4

Typeset by Market House Books Ltd, Aylesbury


Printed in Great Britain by
Butler & Tanner Ltd, Frome

BEST AVAILABLE COPY

Near
G. F.
into
and
was
Dicti
tiona
or m
their
tion
a bio
A
ance
nal l
about
extra
of lit
was
speci
tions
bols,
comm
and /
chem
order
have
terna
so fre
to it
ary, v
reflec
not e
sprea
tional
vario
been
numb
Bio
the st:

igase

Conium maculatum. The natural enantiomer has *S* configuration.



igase γ -Glu-X carboxypeptidase. EC 3.4.19.9 (formerly 3.4.22.12); γ -glutamyl hydrolase; an enzyme that hydrolyzes γ -glutamyl bonds in *N*-pteroyl- γ -oligoglutamate, pteroylglutamate being a prominent product.

igate 1 to effect or undergo **conjugation**. 2 describing a pair of interconvertible substances, e.g. a **conjugate base pair**. 3 the product formed as a result of conjugation.

igate acid-base pair the relationship between two chemical species, BH^+ and B , that are related by the reaction: $BH^+ + H^+ \rightleftharpoons B + H^+$. Then BH^+ is the **conjugate acid** of the base B , B is the **conjugate base** of the acid BH^+ , and BH^+ and B are a conjugate acid-base pair.

igate base see **conjugate acid-base pair**.

igated 1 describing a compound that is formed by the joining of two compounds. 2 displaying or having undergone conjugation.

igated double bond any of two or more double bonds in a molecule where each double bond is separated from the next by one single bond. Compare **conjugation** (def. 2).

igated protein any protein that contains a nonprotein component, often in stoichiometric proportion. The nonprotein component may be a metal ion, a lipid, a carbohydrate, or nucleic acid, and may be either loosely or tightly bound to the polypeptide chain(s).

igation 1 the act of joining together; the state of being joined together. 2 an alternating sequence of multiple and single chemical bonds containing at least two multiple bonds. 3 delocalization of π -electrons and resultant additional structural stability. 3 the covalent or noncovalent joining together of one (larger) molecule, e.g. a protein or bile acid, with a smaller molecule. 4 a process of sexual reproduction occurring in various types of unicellular organisms. In bacteria, e.g. *Escherichia coli*, it involves the transfer of DNA from a donor cell to a recipient cell via a sex pilus; in protozoa, *Paramecium aurelia*, a true exchange of DNA occurs between the participating cells, which belong to different mating strains.

igation labelling a procedure for introducing a label into a molecule of interest by covalently coupling it, in a specific chemical reaction, to a small molecule containing the label (see **conjugation** (def. 3)). It is particularly useful for labelling with radioiodine any protein or peptide that is sensitive to oxidative procedures or to noxious components of commercial radioiodine, that lacks tyrosine residues reactive with iodine, or that requires to be labelled at a residue other than tyrosine. See also **Bolton and Hunter reagent**.

igative plasmid any plasmid that can bring about the transfer of DNA by **conjugation** (def. 3).

igon any genetic element that is required for bacterial conjugation (def. 4), e.g. **fertility factor**.

active tissue any supporting tissue that lies between epithelial tissues and consists of cells embedded in a relatively loose amount of extracellular matrix.

active tissue growth factor *abbr.*: CTGF; a growth factor of testis, spleen, kidney, lung, heart, and brain, that is homologous to **CEP10**; one of a family of growth regulators that belong to a group of immediate-early genes expressed after stimulation by growth factors or certain oncogenes. The family includes **Cyr61**, **β IG-M1**, and **β IG-M2**. Example (precursor) from mouse: database code CTGF_MOUSE, 348 amino acids (35 kDa).


conserved

connectivity (*in chemistry*) the information in any molecular formula or model regarding the order in which the constituent atoms of the molecule are linked, irrespective of the nature of the linkages.

connexin the main protein component of a **connexon**. Each connexin contains four putative membrane-spanning α -helices, and six connexins make up each connexon. A number of different subtypes of connexin exist within each species, contributing different functional behaviour to different connexons. Example from human: database code CX26_HUMAN, 208 amino acids (24.27 kDa); four motifs.

connexon the structural subunit of a **gap junction**, the structure that forms a bridge between adjacent cells in certain tissues of vertebrates. Seen by electron microscopy or low-angle X-ray diffraction, an individual gap junction consists of a number (10^3 – 10^4) of connexons, often in hexagonal array, embedded in and protruding from either side of each of the opposed cell membranes in register and linked to one another. Each connexon is a cylindrical proteinaceous structure, about 6–8.5 nm in diameter and 7.5 nm long; it consists of six rod-shaped, essentially rigid subunits (**connexins**) of about 25 kDa arranged in an annulus. The diameter of the central opening can apparently be varied by radial displacement of the subunits at their cytoplasmic extremities. Each connexin spans one cell membrane, two with connecting ends being required to form a channel across both membranes. Such channels provide a regulable hydrophilic pathway permitting the passage of small molecules (up to about 800 Da) between adjacent cells; the specificity of each channel is determined by the type of connexin it utilizes.

conotoxin *abbr.*: ω -CT; any of several peptides of the family of ω -conotoxins isolated from the venoms of two marine snails, *Conus geographicus* and *C. magus*. All ω -conotoxins have a conserved pattern of cysteine residues linked by four disulfide bridges. They are neurotoxins that inhibit voltage-gated Ca^{2+} channels and neurotransmitter release. ω -Conotoxin GVIA has the structure:



consensus sequence an idealized sequence of nucleotides, or their constituent bases, or amino acids, base, or amino acid that represents the nucleotide most likely to occur at each position in the sequence. Consensus sequences are used to identify RNA splicing sites, other sites, plasmids, and families of proteins.

conservation the retention of structure by a macromolecule, or by a specified segment of one, with variation of circumstance (environmental, genetic, etc.). When used of primary structure, it can be synonymous with **sequence homology**. The degree of retention of structure is usually specified. See also **conserved**.

conservation of energy (law of the) see **thermodynamics**.

conservative base change or **conservative base substitution** any mutational change, including substitution of a base, in a particular base triplet in a DNA molecule such that either the amino acid encoded by that base triplet is not altered, or there is no major change in the properties of the R group of the amino acids involved, e.g. Glu for Asp. See **genetic code**, **wobble hypothesis**.

conservative replacement or **conservative substitution** any replacement or substitution in a polypeptide chain of a particular amino acid residue by another one with similar properties, e.g. Arg for Lys, Phe for Tyr, Glu for Asp.

conserved describing a tendency to invariance in corresponding residues or sequences of residues of encoded macromolecules (e.g. proteins) obtained from specimens of genetically different sources. Macromolecules showing a high degree of